

## METHOD AND SYSTEM FOR TREATING PRESBYOPIA

### CROSS REFERENCE TO RELATED APPLICATIONS

This application claims priority from U.S. provisional application serial no. 60/173,448 filed December 29, 1999; the disclosures of which are incorporated herein by reference.

### BACKGROUND OF THE INVENTION

#### Technical Field

The present invention generally relates to a method and system for performing surgery on the cornea to correct vision defects. More particularly, the present invention relates to a system and method for treating presbyopia by altering the light transmitting properties of the cornea. Specifically, the present invention relates to a system and method for altering the shape of the cornea to have two concentric vision zones that allow a presbyopic patient to focus on near items and far items.

#### Background Information

The human eye transforms light waves into electrical impulses that are interpreted by the brain. Light waves enter the eye through a transparent cornea. The waves then pass through the aqueous humor and through the pupil. The iris expands or constricts the pupil to allow the proper amount of light

to pass to the lens. The light waves then pass through the crystalline lens. The lens is focused with the ciliary muscle so that the light waves are properly focused on the retina. The retina includes light sensitive cells (rods and cones) that transform light waves into electrical impulses. The electrical impulses are transmitted through the optic nerve to the brain. The visual cortex at the back of the brain reconstructs the impulses into an image.

Presbyopia is a natural condition causing the eye to lose its ability to focus on near objects. Over the past several years, presbyopia has been thought to occur due to the loss of flexibility of the crystalline lens in the eye. Presbyopia has traditionally been treated with reading glasses or bifocals that allow the patient to focus on near objects. Surgical correction procedures have not been attempted because of the understanding that presbyopia occurs because the lens has hardened.

In recent years, another theory has developed stating that presbyopia is caused by the growth of the lens causing the ligaments to loosen so that they can no longer exert tension on the lens. A surgical correction method has been developed based on this theory wherein tension is re-introduced to the ligaments supporting the lens by inserting implants.

It is desired in the art to provide a surgical treatment for presbyopia where implants are not required. It is desirable that such a method be as successful and as relatively simple as the widely-used laser refractive surgery methods now known in the art.

### **SUMMARY OF THE INVENTION**

The invention provides a system and a method for treating presbyopia by altering the light transmitting properties of the cornea. The invention provides a method wherein two concentric areas of the cornea are shaped to allow the patient to focus on both near and far objects. The larger area is treated to focus on close objects while the inner smaller area is treated to focus on far objects. The method of the invention may be performed on a system that creates collapsing crescent-shaped eroding laser areas.

The invention allows refractive errors of the eye to be corrected while also correcting the presbyopic condition of the eye.

The invention may also be used to treat myopia.

### **BRIEF DESCRIPTION OF THE DRAWINGS**

Fig. 1 is a cross-sectional view of a typical human eye;

Fig. 2 is a front elevational view of the cornea of the eye;

Fig. 3 is a sectional view taken along line 3-3 of Fig. 2;

Fig. 4A is a view similar to Fig. 2 showing the first stage of a collapsing crescent laser beam applied to the left side of the cornea;

Fig. 4B is a view similar to Fig. 4A showing an intermediate stage;

Fig. 4C is a view similar to Fig. 4A showing a final stage of the collapsing crescent laser beam;

Fig. 5A is a view similar to Fig. 4A showing an initial stage of the collapsing crescent laser beam applied to the right side of the cornea;

Fig. 5B is a view similar to Fig. 5A showing an intermediate stage;

Fig. 5C is a view similar to Fig. 5A showing a final stage of the collapsing crescent laser beam;

Fig. 6A is a view similar to Fig. 4A showing an initial stage of the collapsing crescent laser beam applied to the top half of the cornea;

Fig. 6B is a view similar to Fig. 6A showing an intermediate stage;

Fig. 6C is a view similar to Fig. 6A showing a final stage of the collapsing crescent laser beam;

Fig. 7A is a view similar to Fig. 4A showing an initial stage of the collapsing crescent laser beam applied to the bottom half of the cornea;

Fig. 7B is a view similar to Fig. 7A showing an intermediate stage;

Fig. 7C is a view similar to Fig. 7A showing a final stage of a collapsing crescent laser beam;

Fig. 8 is a sectional view similar to Fig. 3 showing the cornea after the laser treatment of Figs. 4A-7C;

Fig. 9 is a view similar to Fig. 2 showing the cornea of Fig. 8;

Fig. 10 is a sectional view taken along line 10-10 of Fig. 9;

Fig. 11 is a sectional view similar to Fig. 3 showing the resulting shape of the cornea after the treatment of the present invention; and

Fig. 12 is a schematic of the apparatus used to perform the method of the present invention.

Similar numbers refer to similar elements throughout the specification.

## **DESCRIPTION OF THE PREFERRED EMBODIMENTS**

Presbyopia is a condition that affects most human eyes as the eye ages. A typical human eye is depicted in Fig. 1 and is indicated generally by the numeral 10. Eye 10 includes a lens 12 that focuses light waves on the retina 14. The light waves enter the eye through the cornea 16 and pass through the pupil 18 to lens 12. When eye 10 is young, lens 12 can focus light waves on retina 14 from objects that are relatively close to cornea 16. For instance, the young eye may properly focus on the text in a book 20 that is as close as two inches from cornea 16. As the eye ages, lens 12 loses its ability to focus on near objects and eventually book 20 must be held ten to twenty inches away from cornea 16 in order for lens 12 to focus on the text. The young eye thus has an ability to focus in the range indicated by the arrow 22. The old eye 10 can only focus in the range indicated by the numeral 24. This condition is known as presbyopia and has traditionally been treated by supplying reading glasses to supplement lens 12 allowing eye 10 to focus back into range 22. One problem with reading glasses is that the wearer of reading glasses loses his ability to focus on far objects while wearing the reading glasses. The method of the

present invention alters the shape of cornea 16 allowing eye 10 to focus on near objects as well as far objects.

5 The method of the present invention is performed by first determining the shape and focusing ability of eye 10. Various methods for determining these factors are known in the art and any of the variety of methods may be used to gather this information. For instance, the first step may be to use a phoropter to measure the refractive state of eye 10. The person performing the method may optionally determine the topography of cornea 16 with a topographer. The pachymetry or corneal thickness may also be optionally measured.

10 Performing these steps will determine if eye 10 is presbyopic plano, presbyopic with spherical hyperopia, or presbyopic with spherical myopia. It may also be determined that any of these three conditions is combined with astigmatism. When eye 10 has astigmatism, the astigmatism is first treated using a LASIK algorithm to correct for the astigmatism. The patient then must wait for eye 10 to heal (approximately two months) before the method of the present invention is performed.

15 The doctor performing the method next determines an appropriate level of correction that must be applied to cornea 16 to allow eye 10 to focus on objects in range 22. In the first embodiment discussed below, eye 10 is presbyopic but plano meaning that no correction is needed to achieve 20/20 vision other than the correction for the presbyopic condition. In the preferred embodiment of the invention, two diopters of correction will be used to correct

the presbyopic condition. In other embodiments, a different correction may be used. The doctor then selects a central area 30 (Fig. 2) on cornea 16 to perform the initial treatment. In the preferred embodiment, area 30 is a 6mm optical zone.

5 The patient's eye is first anesthetized as is known in the art. The doctor cuts a flap in cornea 16 as is known in the art so that the actual laser treatment on area 30 is performed under the outer surface of cornea 16 in the corneal stroma. In other embodiments, the doctor may remove a disk from the cornea. The disk is replaced after the cornea is sculpted. The Fig. 3 shows the topography of cornea 16 before the method of the present invention is applied to cornea 16 as identified with solid line 32. The initial two diopter treatment will change the shape of cornea 16 to that indicated by dashed line 34. The two diopter treatment steepens the curvature in area 30 so that eye 10 can focus on close objects 20 in range 22.

10 The initial treatment on cornea 16 is performed by eroding area 36 with a laser beam. The boundaries and definition of area 36 is determined using a difference of sphere algorithm or other suitable algorithms known in the art. The resulting area 36 is mathematically defined in a way that a laser can be controlled to remove area 36. The apparatus to sculpt a cornea with a laser beam is known in the art and is preferably a LASIK laser. One example of an apparatus useful for performing this method is disclosed in U.S. Patent 15 5,642,287, the disclosures of which are incorporated herein by reference. The

preferred treatment for removing area 36 is to sculpt cornea 16 with four sequential collapsing crescent-shaped laser beams. In other embodiments of the invention, area 36 may be removed using different known methods such as a scanning spot or an erodible mask. The collapsing crescent-shaped laser beams are applied to area 30 in four steps. Figs. 4A-7C show the application of laser beam 40 to area 30 on cornea 16. The specific sequence of Figs. 4, 5, 6, and 7 is irrelevant to the present invention. The order provided in the drawings is provided as an example but it is understood that the doctor may start on the right half, the upper half, or the lower half instead of the left half as shown in the drawings.

The treatment begins with a half circle-shaped laser applied to the left half of area 30 as depicted in Fig. 4A. The intensity of laser 40 and the amount of time it is applied to area 30 are determined by a controller that analyzes area 36 and determines the time and intensity required for laser 40 to remove area 36. Laser 40 then collapses from the shape shown in Fig. 4A towards the shape shown in Fig. 4C. It is thus understood that the size of laser 40 gradually collapses until it disappears in the direction from the center of area 30 to the edge of area 30.

The doctor then treats the right side of area 30 as depicted in Figs. 5A-5C. Once the left and right sides of area 30 are treated, the doctor changes the laser configuration and treats the top and bottom sections of area 30 as depicted in Figs. 6A-6C and 7A-7C.



The resulting cornea is depicted sectionally in Fig. 8. Treated cornea 16 now has a steepened section 50 that allows eye 10 to focus in range 52 that includes most of range 22, some of range 24, but excludes the far end of range 24. The cornea of Fig. 8 is thus myopic because it cannot focus on objects far away from cornea 16. However, the presbyopic eye can now focus on near objects in range 52.

Having created the cornea Fig. 8, the doctor now selects a smaller area 60 that is substantially concentric with area 30. In the preferred embodiment of the invention, area 60 is a 4mm optical zone. The doctor applies a treatment to area 60 that allows area 60 to focus on objects far away thus treating the myopic condition. This treatment is a spherical diopter correction equal to, but negative of, the positive correction applied to area 30. In this embodiment, a negative 2 diopter correction is performed in area 60.

The negative correction applied to area 60 will remove area 62 of cornea 16 to provide a central area of eye 10 that will be able to focus on far objects. Material 62 is defined and removed using one of the same methods described above.

The resulting cornea is depicted in Fig. 11. Cornea 16 has a central area 70 that is configured to allow eye 10 to focus on far objects and a ring-shaped area 72 that allows eye 10 to focus on near objects. Having undergone this treatment, the presbyopic patient can now focus on near and far objects without the use of implants or reading glasses.

As noted above, the initial examination of eye 10 may determine that eye 10 is hyperopic or myopic. When eye 10 is hyperopic, the correction described above is altered to also correct for the hyperopia. For instance, zone 30 is initially treated with a two diopter correction for the presbyopia along with a correction for the hyperopia. In one example where the hyperopia is plus three diopters, the correction first applied to area 30 is plus three diopters for the hyperopia and plus two diopters for the presbyopia resulting in a positive five diopter spherical correction in zone 30. Zone 60 is then treated with a negative two spherical diopter correction resulting in a correction of the hyperopia in addition to the presbyopia.

When eye 10 is myopic, the initial correction to area 30 includes a combination of the presbyopic correction with a myopic correction. For example, when eye 10 is negative 3 diopters myopic, the presbyopia correction to area 30 is a negative 1 diopter correction in area 30 determined by combining a negative 3 diopter spherical correction with a positive 2 diopter spherical correction. Area 60 is then treated with a negative 2 spherical diopter correction.

After the sculpting operations are performed, a topography of the sculpted eye may be taken to show the difference of curvature in zones 60 and 30. The doctor then examines the patient's distance visual acuity and checks the patient's near visual acuity. If further errors are found, further sculpting is performed.

The apparatus used to perform this method includes a computer 80 that is capable of controlling a laser 82 used to ablate cornea 16. Computer 80 is capable of controlling laser 82 to define and remove areas 36 and 62. Computer 80 may include an input device 84, such as a keyboard, that allows information about eye 10 to be placed in the memory of computer 80. A controller 86 may also be provided that allows computer 80 to communicate with laser 82.

The method and apparatus described above may also be used to treat a myopic cornea. In this treatment, distance is treated in the center and the periphery is treated less to create a solution for the myopic eye. For instance, a 4 millimeter center area may be treated to view distant objects. The peripheral region (outside diameter of 6 millimeters) is treated with a corresponding treatment to allow the eye to see near objects. The 6 millimeter zone is treated first by creating a curvature allowing the 6 millimeter zone to see near objects. A 4 millimeter zone is then treated to flatten the 4 millimeter zone so that it may focus on far objects. The combined zones allow the cornea to focus on near and far objects.

In the foregoing description, certain terms have been used for brevity, clearness, and understanding. No unnecessary limitations are to be implied therefrom beyond the requirement of the prior art because such terms are used for descriptive purposes and are intended to be broadly construed.

Moreover, the description and illustration of the invention is an example and the invention is not limited to the exact details shown or described.

100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000